

Papillary Carcinoma Coding Guideline – Cases dx 2007 to/thru *2017

Rules for Coding Papillary Carcinoma based on Primary Site for Bladder, Thyroid, Breast & Lung

The QC team at the Cancer Registry of Greater California created a Papillary coding Guideline for Bladder, Breast and Thyroid which was distributed regionally. The California Cancer Registry felt this guideline should have a wider distribution to benefit Registrars statewide. Additionally Region 1/8 (CPIC) has contributed one Lung scenario which I have added to the original document. We hope you find this guideline helpful. **Thank you Regional QC Teams!** ~Donna M. Hansen, CTR /CCR Auditor &Trainer

	BLADDER (includes Renal Pelvis, Ureter) - Code 8130				
Histology Text	Code	References			
 Papillary carcinoma Papillary transitional cell carcinoma Papillary urothelial carcinoma Papillary carcinoma and Transitional cell carcinoma 	8130/3	 Renal Pelvis, Ureter, Bladder and Other Urinary Histology Rules, MPH Rule H4 clarifies that these histologies should be coded to 8130 (papillary transitional cell carcinoma). See also, MPH Rules Table 1- Urothelial Tumors SINQ 20130197: Histology code 8050 [papillary carcinoma, NOS] should not be used for papillary carcinoma of the urinary system diagnosed starting in 2007. Rule M6 includes this histology (8050) to take pre-2007 cases into consideration 			
	THYROID – Code 8260 or 8340 depending on if only papillary carcinoma OR papillary & follicular				
Histology Text	Code	References			
Papillary carcinoma	8260/3	 Other Sites Histology Coding Rules, MPH Rule H14 clarifies that if the thyroid tumor is papillary carcinoma, then code papillary adenocarcinoma (8260) SINQ 20071036: For cases diagnosed 2007 or later, assign code 8260 [papillary carcinoma of the thyroid]. 			
Papillary microcarcinoma or micropapillary carcinoma	8260/3	 SINQ 20150023: Per the WHO Tumor of Endocrine Organs, for thyroid primaries/cancer only, the term micropapillary does not refer to a specific histologic type. It means the papillary portion of the tumor is minimal or occult (1cm or less in diameter) and was found incidentally. WHO does not recognize the code 8341 and classifies papillary microcarcinoma of the thyroid as a variant of papillary thyroid and thereby should be coded to 8260. If the primary is thyroid and the pathology states papillary micro-carcinoma or micropapillary carcinoma, code 8260 is correct. 			
Papillary and follicular or papillary with a follicular variant	8340/3	• Other Sites Histology Coding Rules, MPH Rule H15 clarifies that if the thyroid tumor is follicular and papillary carcinoma, then code papillary carcinoma, follicular variant (8340)			
Follicular and papillary microcarcinoma	8340/3	• SINQ 20071076 : For thyroid cancer only, the term micropapillary does not refer to a specific histologic type. It means that the papillary portion of the tumor is minimal or occult, usually < 1 cm in diameter.			



Preventing Cancer. Promoting Life.







"Q"-Tips

May 2016

Quality Tips - Education - Abstracting - Coding

#2016-01

BREAST- Code 8503 when only one histologic type (papillary carcinoma) identified				
Histology Text	Code	References		
Papillary Intraductal carcinoma	8503/2	MPH Rules, Breast Equivalent Terms, Definitions, Tables and Illustrations, Table 1		
Intraductal papillary adenocarcinoma with invasion	8503/3	• MPH Rules, Breast Equivalent Terms, Definitions, Tables and Illustrations, Table 2		
Infiltrating papillary carcinoma	8503/3	 SINQ 20091040: For cases diagnosed 2007 or later, assign histology code 8503 [Papillary adenocarcinoma]. Rule H14 applies. ICD-O-3 code 8050 does not apply in this case. Refer to the numeric listing in ICD-O-3. 8050 is a squamous cell neoplasm. Papillary carcinoma of the breast is NOT a squamous cell neoplasm. It is a neoplasm of the breast parenchyma-ducts, lobules or connective tissue. (Histology code) 8503 is the correct code in this case. 		

LUNG – Code 8260 Micro <i>papillary</i> Adenocarcinoma of lung				
Histology Text	Code	References		
Micropapillary Adenocarcinoma	8260/3	• SINQ 20110115 - An expert pathologist states that WHO notes micropapillary to be a pattern seen in papillary carcinomas, but does not specify it as a separate histologic type.		
		• SINQ 20110125 - The ICD-O-3 codes for micropapillary have specific associations such as ductal, serous or transitional. None of those associations fit lung primaries.		

Content Authors & Contributors				
Papillary Carcinoma Coding Guidelines Bladder, Breast & Thyroid	Micropapillary Adenoca Lung			
Taina Valone, RHIA, CTR	Region 1/8 - CPIC			
Dee LeTendre, RHIT, CTR	Chris Schwarz, BA, CTR			
Winny Roshala, BA, CTR				

Questions? Contact your Regional Quality Control Coordinator					
Region 1	/8 (CPIC)	Shawky Matta, CTR	Shawky.Ma tta@CPIC.org		
	Regions	Dee LeTendre, CTR	dletendre@crgc-cancer.org		
CRGC	2,3,4,6,10				
	Region 5	Crickett Dyke, CTR	<u>ndyke@llu.edu</u>		
Region	9 (CSP)	Mari Carlos, CTR	www.m ylacsp.org		



CANCER PREVENTION INSTITUTE OF CALIFORNIA Preventing Cancer. Promoting Life.



